



Deep Water Aptian Turbidite System in Punta del Este Basin, a New Play Offshore Uruguay

M. Sc. Bruno Conti


Authors: Bruno Conti (1); Santiago Ferro (1); Juan Tomasini (1) Pablo Gristo (1)

1. E&P, ANCAP, Montevideo, Uruguay.



Content

- Importance of turbidites
- Exploration Offshore Uruguay
- Turbidite systems offshore Uruguay
- Aptian-Albian play
- Volumetrics and hydrocarbons evidence
- Conclusions




Content


- **Importance of turbidites**
- Exploration Offshore Uruguay
- Turbidite systems offshore Uruguay
- Aptian-Albian play
- Volumetrics and hydrocarbons evidence
- Conclusions

Importance of turbidites in Atlantic basins

Geological model for gravity flow deposits



The exploratory interest for gravity flow deposits, located in deep marine environments, has grown substantially in the last decades. Recent discoveries made in submarine fans in Atlantic basins of South America (Guyana, French Guiana) and West Africa (Ghana, Liberia, Sierra Leone) encourage the exploration of other frontier areas.






The success of this type of play led to an intense exploration activity in offshore Uruguay (a frontier area with no hydrocarbons discoveries)

These plays:

- Are located in the Atlantic
- In areas with no salt deformation
- In stratigraphic traps
- In Cretaceous fan systems
- In deep waters
- With a Cretaceous source rock



Source Scotese (2014)






Content

- Importance of turbidites
- **Exploration Offshore Uruguay**
- Turbidite systems offshore Uruguay
- Aptian-Albian play
- Volumetrics and hydrocarbons evidence
- Conclusions




Present day seismic data

In the last eight years, as a result of two bidding rounds, an intense exploratory activity was developed offshore Uruguay, including the acquisition of 41,000 km of 2D seismic, 43,600 km² of 3D seismic and the drilling of an ultra-deep exploratory well (after 40 years).



41,000 Km of 2D seismic covering shallow to deep waters





43,600 Km² of 3D seismic covering mostly deep waters



In this context, a great amount of Upper Cretaceous and Cenozoic turbiditic systems were identified with 3D seismic (more than 40 bodies).


- The marine turbidite systems or submarine fans in the offshore basins of Uruguay are mostly located in deep waters and develop in Upper Cretaceous (Conician-Santonian, Maastrichtian) and Cenozoic (Paleocene, Eocene and Oligocene) sequences.
- The turbidites, and associated channels, shows roughly a NW-SE trend, suggesting that the paleoshelf and paleoslope for Upper Cretaceous and Cenozoic times presented an orientation similar to the current margin.
- One Oligocene turbidite was drilled in 2016





Raya well tested an Oligocene turbidite

- WD: 3,404 m
- Sediment depth: 2,452 m
- Reservoir of good quality but water bearing (135 m thick turbidite fan with a 24% average porosity)
- Dry well
- Distance between main source rock and reservoir is around 2,500 m
- 3D seismic show no clear migration pathways between Aptian source rock and this reservoir






Content


- Importance of turbidites
- Exploration Offshore Uruguay
- Turbidite systems offshore Uruguay
- **Aptian-Albian play**
- Volumetrics and hydrocarbons evidence
- Conclusions




3D seismic used in this work (13,000 km²) allowed the interpretation of a Late Aptian marine Fan



Structural map (top horizon)





Amplitude anomaly map (showing fan geometry)







- Despite the great amount of marine turbidites identified for the Upper Cretaceous and Cenozoic, these systems have not been described for the Lower Cretaceous sequences, that mostly show retrogradational stacking patterns.
- However, a big deep water Late Aptian-Early Albian fan system was interpreted






Aptian-Albian Fan in a NW-SE seismic line showing double downlap






This finding is particularly important to better comprehend the evolution of the basin, since the turbidite has a perpendicular orientation (NE-SW) in relation to the younger turbidites, which suggest a different geometry of the basin for that moment






Marine Aptian distribution with SDRs location






Top basement structural map with 2D data



Channels and fans of this age probably restricted to this area of influence of the Rio de la Plata Transfer System



Paleorestitution of the area for the Early K



Creaser et al. (2017)



Detailed top basement structural map of the area with 3D seismic







Content

- Importance of turbidites
- Exploration Offshore Uruguay
- Turbidite systems offshore Uruguay
- Aptian-Albian play
- **Volumetrics and hydrocarbons evidence**
- Conclusions




Volumetric assessment

- Age: Late Aptian - Early Albian
- Bathymetry: 3100 m
- Sediment depth: 5100 m
- Area: 110-486-1337 km²
- Depositional system: Marine fan
- Estimated mean prospective resources: 2364 MMBOE
- Exceeds by far the Minimum Economic Field Size






Prospectivity and hydrocarbon evidences



- ✓ Direct contact with Aptian source rock
- ✓ Inverse polarity with respect to seabed
- ✓ Abrupt increase in seismic amplitude (bright spot)
- ✓ AVO anomalies associated with the fan
- ✓ Gas chimneys coming from Aptian source rock feeding Upper cretaceous reservoirs





Content

- Importance of turbidites
- Exploration Offshore Uruguay
- Turbidite systems offshore Uruguay
- Aptian-Albian play
- Volumetrics and hydrocarbons evidence
- **Conclusions**

- More than 40 Upper Cretaceous and Cenozoic turbidite systems were identified offshore Uruguay in the last 10 years.
- In this work 3D seismic analysis allowed the identification of a deep water Aptian turbidite system in a transition area between Punta del Este and Pelotas basins.
- It has a perpendicular trend with respect to the Upper K and Cenozoic turbidites. The later were generated in a passive margin phase, while the Aptian Fan System was developed in a transitional phase with a different geometry of the basin
- Provenience of sediments are from the Rio de la Plata craton (good reservoir quality is expected)
- There are DHI on the seismic that confirm the prospectivity of the prospect (amplitude anomalies, AVO anomalies, inverse polarity, gas chimneys).
- The fan is in direct contact with the Aptian source rock, which is a world class source rock, reducing migration risks.

These conclusions make offshore Uruguay attractive and also opens a new play type for hydrocarbon exploration.



Acknowledgment

The authors would like to thank ANCAP for the data provided for this investigation.

References

- **Creaser, A.; Hernández-Molina, F.; Badalini, G.; Thompson, P.; Walker, R.; Soto, M.; Conti, B.** 2017. A Late Cretaceous mixed (turbidite-contourite) system along the Uruguayan Margin: Sedimentary and palaeoceanographic implications. *Marine Geology*. p.234-253.
- **Scotese, C.R., 2014.** Atlas of Neogene Paleogeographic Maps (Mollweide Projection), Maps 1-7, Volume 1, The Cenozoic, PALEOMAP Atlas for ArcGIS, PALEOMAP Project, Evanston, IL.
- **Sempstrata. 2019.** <http://www.sepmstrata.org/page.aspx?pageid=259> . Accessed march 1st, 2019
- **Soto, M.; Morales, E.; Veroslavsky, G.; De Santa Ana, H.; Ucha, N. & Rodriguez, P. 2011.** The continental margin of Uruguay: Crustal architecture and segmentation. *Marine and Petroleum Geology*, Volume 28, Issue 9, September 2011, p. 1676-1689.